

Energy expenditure and nutrition status of ballet, jazz and contemporary dance students

Rossioud D.^{1 A,B,D-F}, Papadopoulou S.^{1 B,D-F}, Pagkalos I.^{2 A,C,D,F}, Kokkinopoulou A.^{1 B,D,F}, Petridis D.^{3 C,D,F}, Hassapidou M.^{1 A,D,F*}

1. Department of Nutrition & Dietetics, Alexander Technological Educational Institute of Thessaloniki. Sindos, Thessaloniki. Greece
2. Department of Electrical & Computer Engineering, Aristotle University of Thessaloniki. Thessaloniki, Greece
3. Department of Food Technology, Alexander Technological Educational Institute of Thessaloniki, Sindos, Thessaloniki. Greece

A - Conception and study design; **B** - Collection of data; **C** - Data analysis; **D** - Writing the paper; **E** - Review article; **F** - Approval of the final version of the article; **G** - Other (please specify)

ABSTRACT

Purpose: To evaluate of the energy expenditure in 3 types of dance classes (ballet, jazz and contemporary), as well as of the daily energy balance depending on the type of dance.

Materials and methods: 40 females attending dance classes with a median age of 21.0 (19.0-25.0) and 10 males with a median age of 27.0 (20.0-28.0) participated in this study. The energy cost of each dance class was measured using the BodyMedia SenseWear Sensor and total daily energy expenditure was evaluated using a 3-day recording of physical activity. The dietary intake was evaluated with a 3-day food diary recording. Statistical analysis was performed using the SPSS software.

Results: Median energy expenditure varied from 306 (277-328) Kcals/class for contemporary dance to 327 (290-355) Kcals/class for ballet and 369 (333-394) Kcals/class for jazz for females with

significant differences between contemporary and jazz classes. For males energy expenditure was 508 (447-589) Kcals/class and 564 (538-593) Kcals/class for ballet and jazz classes, respectively. Females had lower values for all anthropometric measurements, energy intake, macronutrient intakes and energy expenditure, comparing to males. The anthropometric characteristics did not differ between types of dance. Both female and male dance students were in negative energy balance.

Conclusions: The use of sensors such as Bodymedia Sensewear together with keeping daily diaries make measurement of physical activity in dancing reliable and accurate. Exercise expenditure differs across types of dance in females but not in males. Both gender had inadequate energy and carbohydrate intakes.

Key words: Energy balance, dietary intake, BodyMedia SenseWear, dance

*Corresponding author:

Hassapidou Maria
Department of Nutrition & Dietetics
Alexander Technological Educational Institute of Thessaloniki
Thessaloniki, Greece
Tel.: +302310013591. e-mail: mnhas@nutr.teithe.gr

Received: 08.01.2017

Accepted: 08.03..2017

Progress in Health Sciences

Vol. 7(1) 2017 pp 7-17

© Medical University of Białystok, Poland

INTRODUCTION

The World Health Organization (WHO) defined physical activity (PA) as “bodily movement produced by skeletal muscles that requires energy expenditure”, while physical inactivity reaches the fourth place as a risk factor for global mortality [1]. A large number of methods with differences in validity and reliability, can be used for PA assessment. with the two most used methods being self-report questionnaires and wearable sensors. such as accelerometers [2]. It is well established that PA contributes to a healthy lifestyle. by preventing and modifying chronic diseases, like obesity, metabolic syndrome, type II diabetes, cardiovascular diseases and some types of cancer [3].

Aerobic dance consists one of the most favoured adult fitness activities, especially for women, and it could be a cardiovascular training alternative, if performed according to the guidelines of the American College of Sports Medicine (ACSM) [4]. The amount of energy expenditure during a bout of aerobic dance varies according to the intensity of the exercise and style, as well. Thus, dance exercise may require a variety of cost from 4 to 11 kcal/minute [5]. Various forms of dance found to be as effective as traditional modalities. such as jogging. meeting the ACSM guidelines for weight modification and maintenance.

Energy expenditure measurement in order to assess energy balance as well as fuel utilisation and thermogenesis during exercise is required [6]. According to Levine et al. [7] there was a significant thermogenic potential of fidgeting-like at low workloads. which contributes substantially to energy balance. Energy expenditure increased while sitting without motion by 4%, while fidgeting while seated by 54%. while standing motionless by 13%, while fidgeting while standing by 94%, compared to metabolic rate in the supine position.

Two categories of physical activity assessing methods are widely used; subjective (i.e. physical activity questionnaires and diaries) and objective methods (i.e. measures of energy expenditure, physiologic measures, motion sensors). Nowadays, technological advances can provide precision in measures of energy expenditure in free-living individuals by using wearable devices. In other words, wearable sensors give information about the total amount, frequency, duration, and intensity of physical movement [8]. Tri-axial accelerometers detect body displacement electronically in three axes and consist an objective and reliable method to measure the intensity of movement and physical activity pattern. Thus, data obtained using tri-axial accelerometers were well correlated with the total daily energy expenditure. measured using doubly labelled water, divided by basal metabolic rate (BMR) [9].

There is a paucity of data regarding anthropometric measurements energy cost of dancing and energy balance assessment in dance students and to the best of our knowledge there is no such study in Greece. Thus. the purpose of our study was to assess (i) energy expenditure in three types of dance (ballet, jazz and contemporary) using wearable sensors and physical activity records. (ii) energy intake using food diaries. For all the participating individuals a number of anthropometric measurements were recorded.

MATERIALS AND METHODS

Population sample

This is a random study. Dancers volunteered to participate in the study. responding to a posted announcement suspended in a dance school in Thessaloniki. Greece. Participants were informed about the protocols study. All athletes were instructed to follow their usual activity and nutrition dairy prior to the study. Forty female dancers (8 adolescents and 32 adults) attending dance classes for more than two years (14 jazz, 16 ballet and 10 contemporary dancers). and 10 adult males (6 jazz. 4 ballet dancers) with median age of 21 (19.0-25.0) years and 27 (20.0-28.0) years. respectively participated in this study. The study was carried out in Thessaloniki, Greece, in 2014.

All participants have given their written consent prior to the effectuation of the study. which received approval by the Ethics Committee of the Alexander Technological Educational Institute. in Thessaloniki. Greece.

Anthropometric measurements

An experienced dietician performed all anthropometric measurements in the morning, with participants in minimal clothing. Height was measured to the nearest 0.5 cm using a stadiometer, with an accuracy of 0.5 cm (SECA 220, Seca Corporation. Columbia. USA). Body weight was measured with the use of a calibrated digital scale with an accuracy of ± 100 g (Seca 707, Seca Corporation, Columbia, USA). Body Mass Index (BMI) was calculated as body weight divided by the square of height (kg/m^2). Waist circumference was measured with a tape midway between the top of the iliac crest and the bottom of the rib cage. at the end of gentle expiration. Hip circumference was measured with a tape over the great trochanters. Both circumferences were measured over the naked skin. Waist to Hip Ratio (WHR) was defined as waist circumference (cm)/hip circumference (cm), with results ≥ 1.0 and ≥ 0.85 indicating obesity for males and females respectively. according to WHO [10].

Physical activity

Assessment of physical activity was performed using wearable sensors and physical activity records. Energy cost of each dance session (ballet, jazz and contemporary) was measured with the BodyMedia SenseWear Sensor and the records were analyzed in SenseWear Software 7.0. Furthermore total daily energy expenditure was estimated for three consecutive days (two weekdays and one weekend day) from completed physical activity records for the same days. Ainsworth et al. [11] tables were used to estimate the energy cost for each activity. Basal metabolic rate was estimated with the use of Harris & Benedict equations [12].

Dietary intake

All subjects kept a three-day food diary, which consisted of two weekdays and one weekend day, in order to reflect valid results and typical intake [13]. The days of assessing dietary intake and physical activity were the same. A registered dietitian provided verbal and written instructions on how to complete their food diaries. The use of food models helped the subjects estimate the quantity of foods. A computerized nutrient analysis program “Food Processor” (version 7.4) with Greek foods included was used for the analysis of food records. Macronutrient intake was expressed as a percentage of the Recommended Dietary Allowance (RDA) [14] in order to assess the adequacy of micronutrient intake.

Statistical analysis

Statistical analysis was performed using SPSS version 17.0 (SPS Inc., Chicago, IL, USA). Due to the small sampling size non parametric tests were applied. Mann-Whitney U tests for two independent samples were performed for comparison among the two sexes. In order to examine differences between the three types of dance Kruskal-Wallis ANOVA by ranks was used. Results are presented as median in joint with the 1st and 3rd quartile (corresponding to 25% and 75% of data). Dancers’ life style (profession, cooking, dairy activity, marital status etc.) was investigated in concert with gender effect using cross-tabulated information and chi-square statistical analysis. Statistical significance was set at $p < 0.05$.

RESULTS

Age distribution of male and female dancers is presented in Figure 1. 16% of the sample were adolescents and all of them females.

Female dancers had statistically lower median values for weight, height, BMI (Body Mass Index), wrist, waist circumferences, WHR ($p < 0.01$) and hip circumference ($p < 0.05$) when comparing to male dancers (Table 1).




Figure 1. Age distribution of male and female dancers. Females are positively skewed and males negatively

Table 1. Anthropometric characteristics. according to gender (median and quartiles 1st and 3rd)

	Men n=10	Women n=40
Age (yrs)	27.0 (20.0-28.0)	21.0 (19.0-25.0)
Weight (Kg)	78.0 (61.70-81.60)**	55.80 (49.90-59.45)**
Height (cm)	185.0 (173.0-185.50)**	165.0 (160.0-169.0)**
BMI (Kg/m²)	22.79 (20.62-23.99)**	19.77 (18.92-20.85)**
Wrist circumference (cm)	18.0 (15.0- 18.5)**	14.75 (14.0-15.0)**
Waist circumference (cm)	89.0 (79.0-90.0)**	70.50 (67.50-75.25)**
Hip circumference (cm)	101.0 (91.0-102.0)*	91.0 (89.0-96.50)*
Waist-to-hip ratio	0.87 (0.84-0.88)**	0.76 (0.75-0.81)**

*p <0.05. **p<0.01

As shown in Table 2, female dancers had also. statistically lower median BMR, (p<0.01) than male dancers. Both male and female dancers had less energy intake (EI) than their energy expenditure, resulting in negative energy balance, with males

having higher negative values than females (p<0.05). Male dancers had statistically significant higher values in energy intake and higher consumption in grams of protein, fat and dietary fiber, than females (p<0.05).

Table 2. Energy and macronutrient intake and energy expenditure. according to gender (median and quartiles 1st and 3rd)

	Men n=10	Women n=40
Basal Metabolic rate (Kcal)	1882 (1645-1882)**	1376 (1330-1429)**
Energy Intake (EI) (Kcal)	2297 (1703-2374)*	1708 (1381-2097)*
Energy Balance (Kcal)	-1115 (-2153 - -647)*	-526 (-857 - -190)*
Protein (g)	131.24 (54.01-146.40)*	68.89 (49.36-82.0)*
Protein (g/Kg BW)	1.68 (0.88-1.69)	1.19 (0.97-1.64)
Protein (%EI)	21.16 (9.10-30.83)	14.81 (12.86-17.89)
Carbohydrates (g)	244.97 (146.90-324.02)	209.08 (156.90-261.67)
Carbohydrates (g/Kg BW)	3.56 (1.88-5.25)	3.51 (2.80-5.40)
Carbohydrates (%EI)	41.04 (34.51-54.59)	46.78 (41.25-57.62)
Fat (g)	108.09 (68.41-127.97)*	71.21 (60.76-85.37)*
Fat (g/Kg BW)	1.52 (0.88-1.75)	1.31 (1.09-1.64)
Fat (%EI)	36.16 (36.16-40.97)	39.16 (29.46-43.25)
Saturated Fat (g)	31.33 (25.25-41.11)	24.53 (18.69-38.72)
Alcohol (g)	0.0 (0.0-0.0)	0.0 (0.0-0.0)
Dietary Fiber (g)	21.30 (19.23-23.36)*	16.98 (12.05-20.30)*

*p<0.05. **p<0.01

According to Table 3, female dancers had statistically higher median intake in fruits (p<0.01) and milk (p<0.05), while male dancers had higher

median consumption in vegetables, meat and meat products (p<0.05), and fat (p<0.01).

Table 3. Food groups servings intake. according to gender (median and quartiles 1st and 3rd)

	Men n=10	Women n=40
Bread and cereal (servings/day)	6.55 (4.90-8.20)	6.25 (4.20-7.75)
Vegetables (servings/day)	5.20 (3.30-6.70)*	2.25 (0.90-3.05)*
Fruits (servings/day)	0.0 (0.0-0.0)**	1.80 (0.95-3.50)**
Meat and products (servings/day)	5.05 (0.80-5.10)*	1.10 (0.70-2.40)*
Milk (servings/day)	0.25 (0.0-0.50)*	1.95 (1.20-3.40)*
Added Fats (servings/day)	18.05 (14.50-21.60)**	7.20 (3.95-12.20)**

*p<0.05, **p<0.01

The analysis between the three types of dances was performed independently for the two genders and is shown in Table 4 and 5. Female ballet dancers were younger than contemporary dancers, whereas female jazz dancers had higher median exercise expenditure

(in Kcal) than contemporary dancers ($p < 0.05$). Male ballet dancers had no significant differences in anthropometric characteristics or energy expenditure according to type of dance (Table 5).

Table 4. Anthropometric characteristics of female dancers according to type of dance (median and quartiles 1st and 3rd). Different letters denote statistically significant differences among types of dance

	Jazz n=14	Ballet n=16	Contemporary n=10
Age (yrs)	21.0 (19.0-30.0)* ^{ab}	20.50 (15.0-21.0)* ^b	24.0 (21.0- 31.0)* ^a
Weight (Kg)	54.50 (49.90-57.60)	55.80 (50.60-60.80)	53.40 (46.70-58.0)
Height (cm)	163 (160-168)	166.5 (164-170)	162.50 (160.0-170.0)
BMI (Kg/m²)	19.90 (19.0-21.70)	20.0 (19.40-20.70)	19.60 (18.20-19.90)
Wrist circumference (cm)	15.0 (14.0-15.0)	14.80 (13.80-15.0)	14.30 (14.0-15.0)
Waist circumference (cm)	71.50 (68.0-76.0)	72.50 (68.50-76.30)	69.0 (64.0-70.0)
Hip circumference (cm)	91.0 (89.0-97.0)	94.0 (89.50-97.50)	89.50 (85.0-95.0)
Waist-to-hip ratio	0.78 (0.75-0.83)	0.77 (0.75-0.80)	0.75 (0.72-0.81)
Basal Metabolic rate (Kcal)	1371 (1330-1401)	1427 (1337-1459)	1341 (1290-1380)
Energy expenditure (Kcal)/class	369 (333-394)* ^a	327 (290-355)* ^{ab}	306 (277-328)* ^b
Mets/class	4.20 (3.90-4.60)	3.70 (3.40-4.40)	3.70 (3.50-4.20)

* $p < 0.05$

Table 5. Anthropometric characteristics of male dancers according to type of dance (median and quartiles 1st and 3rd)

	Jazz n=6	Ballet n=4
Age (yrs)	27.0 (20.0-28.0)	27.0 (3.50-29.0)
Weight (Kg)	78.0 (61.70-84.80)	78.0 (69.90-79.80)
Height (cm)	185.0 (173.0-188.0)	182.50 (176.50-185.0)
BMI (Kg/m²)	22.80 (20.60-24.0)	22.80 (21.70-24.0)
Wrist circumference (cm)	18.0 (15.0-18.0)	17.0 (15.50-18.0)
Waist circumference (cm)	89.0 (79.0-89.50)	86.0 (81.0-89.50)
Hip circumference (cm)	101.0 (91.0-106.0)	100.0 (95.0-101.0)
Waist-to-hip ratio	0.87 (0.84-0.88)	0.88 (0.86-0.89)
Basal Metabolic rate (Kcal)	1882 (1645-1983)	1881 (1762-1882)
Energy expenditure (Kcal)/class	564 (538-593)	508 (447-589)
Mets/class	4.90 (4.60-5.10)	4.85 (4.25-5.05)

According to dancers' shopping and cooking habits, female dancers reported that their grandma (2.5%), their mother (55.0%) and themselves (42.5%) cooked, while all male dancers reported that their mother did the cooking. Female dancers reported that their grandma (2.5%), their mother (52.5%) and themselves (45.0%) did the shopping, while the respective values for male dancers were 10.0%, 70.0% and 20.0%. According to their profession, female dancers reported that they were self-employed workers (10.0%), teachers (15.0%) and students (75.0%), while the respective values for male dancers were 50.0%, 0.0% and 50.0% ($p < 0.05$). Regarding their

dairy activity, female dancers revealed that 70.0% worked up to 8h/d and 30.0% above 8h/d. with males having respective values of 60.0% and 40.0%. Female dancers spent idle hours up to 5h/d at 60.0% and above 5h/d at 40%. while the respective values for men were 40.0% and 60.0%. Sleeping and physical activity distributed differently according to gender. Thus, 70% of female slept up to 8h/d and 30.0% above 8h/d. whereas all males slept up to 8h/d ($p < 0.05$). The overwhelming majority of female dancers (82.50%) spent up to 3h/d and 17.5% above 3h/d in vigorous activities. with respective values in men 30.0% and 70.0% ($p = 0.01$).

DISCUSSION

The present study revealed that female dancers had lower median values for weight, height, BMI, wrist, waist and hip circumferences, waist to hip ratio, body size, and basal metabolic rate compared to male dancers. Stensland and Sobal [15] and Yannakoulia and colleagues [16] found similar anthropometric characteristics for all three types of dances. In a study conducted in the US and Greece respectively. Also, the dancers of the present study were taller and heavier than female professional ballet dancers from Kuala Lumpur [17], while they had similar BMI values. American professional ballet dancers [18] had similar weight, height, and BMI with the dancers of the present study. Furthermore, similar BMI values are found in elite adolescent female figure skaters in the USA [19] and in dancers from the US National Figure Skating Dance Team [20]. Most of the studies conclude that low body weight and BMI are requisite in any dancing movement valued for both aesthetic and mechanical reasons [17,20].

Female Greek dancers of the study had statistically lower exercise expenditure during dance classes than male dancers. What is more, comparisons according to gender showed that females' energy expenditure was higher in jazz class compared contemporary dance class, while in males there was no difference in energy expenditure according to type of dance.

Water consumption was insufficient in both females and male dancers of the study, while female dancers drank only 0.72lt and male dancers 1.25lt during an exercise session of 1.5 hours. Athletes do not voluntarily drink adequate water to prevent dehydration during exercise, and this can provoke dehydration that compromise athletic performance. Replacement of fluids should compensate sweat and urine losses and this requires generally 200 mL to 300 mL every 15 minutes [21].

Energy intake and energy balance

Greek female dancers' intake (1708Kcal/d. 1381-2097) was quite lower than the energy intake of female ballet dancers in Krakow according to Zulawa and colleagues [22] (1965.48Kcal/d) and slightly higher when comparing to a different Greek dancers female group as reported from Yannakoulia et al. [18]. Also, a group of Spanish ballet dancers [23] consumed a hypocaloric diet of 1555Kcal/d and USA female figure skaters had reports of 1416Kcal/d and 1491Kcal/d when examined by Ziegler et al. [20] and Dwyer et al. [19] respectively. As far as Greek male dancers are concerned, they had a median energy intake of

2297Kcal/d (1703-2374), while US male figure skaters appeared to have a higher energy intake (2837Kcal/d) [20].

Both Greek male and female dancers had statistically lower energy intake than energy expenditure, leading to a negative energy balance e-1115 (-2153 --647) Kcal/d for males and -526 (-857- 190) Kcal/d for females, with males having higher negative values than females. Similarly, a negative energy balance has been demonstrated in former Greek studies [24]–[26]. Large discrepancies between energy intake and energy requirement of athletes of stable body weight can be attributed to underreporting or overestimation of energy needs in days with no exercise [19].

Macronutrient intake

In the present study, the median daily carbohydrate intake was 3.56 (1.88-5.25) g/Kg BW for men and 3.51 (2.80-5.40) g/Kg BW for women. These values are much lower than the current carbohydrate recommendation for athletes (6-10g/Kg BW/day) [27] which could provoke depletion in glycogen stores over time [28]. Female dancers of the present study had lower carbohydrate consumption than adolescent female skaters (4.8g/Kg BW) [19]. Female Greek dancers of Yannakoulia et al. consumed inadequate carbohydrates with reports of 47.9% EI [16]. Similarly, carbohydrate intake of both male and female US figure skating dancers was also insufficient (44% EI and 51% EI, respectively) [20].

Median protein intake of Greek dancers was 1.68 (0.88-1.69) g/Kg BW for males and 1.19 (0.97-1.64) g/Kg BW for females, values that met the respective recommendations for athletes (1.2-1.4g/Kg BW/day) [29]. Regarding fat intake, dancers in the study consumed fat-dense diet with a median fat contribution of 36.16 (36.16-40.97) % to EI for men and 39.16 (29.46-43.25) % EI for women, respectively, values that are higher than the recommendations [27]. Female dancers of the present study had higher protein intake compared to young female dancers of Yannakoulia et al. (1g/kg BW or 15.4% EI, while both groups exhibited high fat consumption [16]. Also, Spanish ballet dancers consumed a high density diet [25]. Protein and fat intake of both male and female US figure skating dancers were higher than the recommendations, in the study of Ziegler et al. [20], while according to a study conducted 11 years later, by Dwyer et al. [19], USA adolescent female skaters showed a lower fat intake (23.8% EI) and a similar protein intake (1.2 g/Kg BW) compared to Greek female dancers.

Male dancers in the present study had significantly higher consumption in protein, fat and also in dietary fiber than female dancers. Higher protein and fat consumption by males can partly explained by the fact that male dancers had higher consumption in meat and fat food group equivalents compared to females. Female dancers had also higher intake in fruits and milk than males.

Greek male dancers were more active during the day compared to females as males had significantly fewer sleeping hours and significantly more hours for vigorous activity than women. Males worked as freelancers in significant higher percentage compared to females. The use of wearable sensors together with keeping daily diaries make measurement of physical activity reliable and accurate. Dancing is a high energy cost exercise that should be recommended particularly in adolescents and young adults, because it contributes significantly to an increase in energy expenditure and a healthy body weight.

Future studies are needed (i) to further examine the effects of nutritional status on performance of the dancers, (ii) to inform the training staff and the athletes on nutritional guidelines and (iii) to develop strategies to improve their dietary intake.

Strengths and limitations

The biggest strength of the study is the use of wearable sensors, which give information about the total amount, frequency, duration, and intensity of physical movement. They are capable to detect small movements, can be used with people of all ages, and in general are more accurate comparing to self-reported PA instruments. The Bodymedia SenseWear Sensor should be added to the strengths of the study, as it is an accelerometer clinically validated and appears in more than hundred papers regarding energy expenditure evaluation in humans [30]. Therefore the results of this study on the energy cost of different types of dancing are reliable and accurate.

The self-reported food diaries that were used for the assessment of dietary intake are inexpensive, and easily applicable, but the danger of underreporting is a well-known limitation.

With the exception of one study [15], the participants had similar anthropometric and nutritional characteristics with other published studies on dancers [16-20,22,23], although the analysis and classification of the intake in food groups has only been seen in this study.

On the other hand, conclusions of the results cannot be generalized for the whole country, as participants were from the same city

(Thessaloniki, Greece) and from the same dance school.

CONCLUSIONS

In view of the results of this study, the use of sensors such as Bodymedia Sensewear together with keeping daily diaries make measurement of physical activity in dancing reliable and accurate. Female jazz dancers had higher exercise expenditure (in Kcals) than contemporary dancers ($p < 0.05$), while male ballet dancers had no significant differences in energy expenditure according to type of dance. Both male and female dancers had a negative energy balance and inadequate carbohydrate intake. In order to compensate the high-energy demands, their intakes during the exercise days should be increased.

Conflicts of interest

None disclosed.

Funding

Supported by a grant of NSRF 2007-2013.

REFERENCES

1. World Health Organization. Global recommendations on Physical Activity for health. World Health Organization; 2010.
2. Westerterp KR. Assessment of physical activity: a critical appraisal. *Eur J Appl Physiol.* 2009 Apr;105 (6):823-8.
3. Papadopoulou SK, Papadopoulou SD, Zerva A, Paraskevas GP, Dalkiranis A, Ioannou I, Fahantidou A. Health status and socioeconomic factors as determinants of physical activity level in the elderly. *Med Sci Monit.* 2003 Feb;9(2):CR79-83.
4. Pollock ML, Gaesser GA, Butcher JD, Després JP, Dishman RK, Franklin BA, Garber CE. (1998). ACSM position stand: the recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults. *Med Sci Sports Exerc.* 1998 Jun;30(6):975-91.
5. Williford DHN, Scharff-Olson M, Blessing DL. The physiological effects of aerobic dance. *Sports Med.* 2012 Dec;8(6):335-45.
6. Levine JA. Measurement of energy expenditure. *Public Health Nutr.* 2005 Oct; 8(7A):1123-32.
7. Levine JA, Baukol PA, Westerterp KR. Validation of the Tracmor triaxial accelerometer system for walking. *Med. Sci. Sports Exerc.* 2001 Sept 33(9):593-7.

8. Strath SJ, Kaminsky LA, Ainsworth BE, Ekelund U, Freedson PS, Gary RA, Richardson CR, Smith DT, Swartz AM. Guide to the assessment of physical activity: clinical and research applications. *Circulation*. 2013 Nov;128(20):2259-79.
9. Bouten CV, De Venne WPV, Westerterp KR, Verduin M, Janssen JD. Daily physical activity assessment: comparison between movement registration and doubly labeled water. *J Appl Physiol*. 1996 Jan;81(2):1019-26.
10. Consultation, WHO Expert. Waist circumference and waist-hip ratio. Report of a WHO Expert Consultation Geneva: World Health Organization, 2008, 8-11.
11. Ainsworth B, Haskell W, Whitt M, Irwin ML, Swartz A, Strath S, O'Brien W, Bassett D, Schmitz K, Emplaineourt P. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc*, 2000; 32:S498-S504.
12. Frankenfield DC, Muth ER, Rowe WA. The Harris-Benedict Studies of Human Basal Metabolism. *J Am Diet Assoc*. 1998 Apr; 98(4):439-45.
13. Crawford PB, Obarzanek E, Morrison J, Sabry ZI. Comparative advantage of 3-day food records over 24-hour recall and 5-day food frequency validated by observation of 9- and 10-year-old girls. *J Am Diet Assoc*. 1994 Jun;94(6):626-30.
14. Washington DC. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients). National Academies Press; 2005.
15. Stensland SH, Sobal J. Dietary practices of ballet, jazz, and modern dancers. *J Am Diet Assoc*. 1992 Mar;92(3):319-24.
16. Yannakoulia M, Sitara M, Matalas AL. Reported eating behavior and attitudes improvement after a nutrition intervention program in a group of young female dancers. *Int J Sport Nutr Exerc Metab*. 2002 Mar;12(1):24-32.
17. Hidayah GN, Bariah AHS. Eating attitude, body image, body composition and dieting behaviour among dancers. *Asian J Clin Nutr*. 2011;3(3):92-102.
18. Kaufman BA. Bone Density and Amenorrhea in Ballet Dancers Are Related to a Decreased Resting Metabolic Rate and Lower Leptin Levels. *J Clin Endocrinol Metab*. 2002 Jun;87(6):2777-83.
19. Dwyer J, Eisenberg A, Prelack K, Song WO, Sonnevile K, Ziegler P. Eating attitudes and food intakes of elite adolescent female figure skaters: a cross sectional study. *J Int Soc Sports Nutr*. 2012;9(1):53.
20. Ziegler PJ, Jonnalagadda SS, Lawrence C. Dietary intake of elite figure skating dancers. *Nutr Res*. 2001;21(7):983-92.
21. Casa DJ, Armstrong LE, Hillman SK, Montain SJ, Reiff RV, Rich BSE, Roberts WO, Stone JA. National Athletic Trainers' Association position statement: fluid replacement for athletes. *J Athl Train*. 2000 Apr-Jun;35(2):212-24.
22. Zuława G, Pilch W. The estimation of nutrition habit of ballet school students in Krakow. *Rocz Państw Zakładu Hig*. 2012;63(1):105-10.
23. Loucks AB. Energy balance and body composition in sports and exercise. *J Sports Sci*. 2004 Jan;22(1):1-14.
24. Papadopoulou SK, Gouvianaki A, Grammatikopoulou MG, Maraki Z, Pagkalos IG, Malliaropoulos N, Hassapidou MN, Maffulli N. Body composition and dietary Intake of elite cross-country skiers members of the Greek national team. *Asian J Sports Med*. 2012 Dec;3(4):257-66.
25. Papadopoulou SK, Papadopoulou SD. Comparison of nutritional intake between volleyball and basketball women athletes of the olympic national teams. *Gazz Med Ital Arch Sci Med*. 2008;167(4):147-52.
26. Papadopoulou SK, Papadopoulou SD, Gallos GK. Macro- and micro-nutrient intake of adolescent Greek female volleyball players. *Int J Sport Nutr Exerc Metab*. 2002;12(1):73-80.
27. Rodriguez NR, DiMarco NM, Langley S. Position of the American dietetic association, dietitians of Canada, and the American college of sports medicine: nutrition and athletic performance. *J Am Diet Assoc*. 2009 Mar;109(3):509-27.
28. Calder PC, Jackson AA. Undernutrition, infection and immune function. *Nutr Res Rev*. 2000 Jun;13(1):3-29.
29. Nogueira JAD, Da Costa THM. Nutritional status of endurance athletes: what is the available information?. *Arch Latinoam Nutr*. 2005;55(1):15-22.
30. <http://sensewear.bodymedia.com/SenseWear-Studies/SW-Bibliography>. [cited 2017 Jan 20].